

Calm and Frenzy: marine obligate hydrocarbonoclastic bacteria sustain ocean wellness

Yakimov, Michail M.; Bargiela, Rafael; Golyshin, Peter

Current Opinion in Biotechnology

DOI:

[10.1016/j.copbio.2021.09.015](https://doi.org/10.1016/j.copbio.2021.09.015)

Published: 02/02/2022

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Yakimov, M. M., Bargiela, R., & Golyshin, P. (2022). Calm and Frenzy: marine obligate hydrocarbonoclastic bacteria sustain ocean wellness. *Current Opinion in Biotechnology*, 73, 337-345. <https://doi.org/10.1016/j.copbio.2021.09.015>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Calm and Frenzy: marine obligate hydrocarbonoclastic bacteria sustain**
2 **ocean wellness.**

3 Michail M Yakimov¹, Rafael Bargiela² and Peter N Golyshin²

4

5 Addresses:

6 ¹Institute of Polar Sciences, Italian National Research Council, 98122 Messina, Italy

7 ²Centre for Environmental Biotechnology, School of Natural Sciences, Bangor University,
8 LL57 2UW Bangor, United Kingdom

9

10 Corresponding author: Yakimov, Michail M (mikhail.iakimov@cnr.it)

11

12

13 **Abstract**

14 According to current estimates, the annual volume of crude oil entering the ocean due to
15 both anthropogenic activities and naturally-occurring seepages reaches approximately 8.3
16 million metric tons. Large discharges from accidents such as with tankers '*Amoco Cadiz*'
17 and *M/T 'Haven'* and with the oil rig '*Deepwater Horizon*' have caused large-scale
18 environmental disasters with extensive damage to the marine ecosystem. Typically, oil
19 cleaning operations take months (and sometimes years) to bring the areas around an
20 accident back to normality. The natural clean-up of petroleum spills in marine environments
21 is carried out primarily by naturally-occurring, taxonomically diverse, obligate
22 hydrocarbonoclastic bacteria (OHCB) that are capable of using hydrocarbons as a sole
23 source of carbon and energy. The natural hosts of OHCB include a range of marine primary
24 producers, e.g. unicellular photosynthetic eukaryotes and cyanobacteria, which have been
25 documented as both, suppliers of hydrocarbon-like compounds that fuel the 'cryptic'
26 hydrocarbon cycle (estimated to 100-500-fold more hydrocarbon than technogenic sources)
27 and as a source of isolation of new OHCB. A very new body of evidence suggests that
28 OHCB are not only the active early-stage colonizers of plastics and hence an important
29 component of the ocean's 'plastisphere' but also encode an array of enzymes
30 experimentally proven to act on petrochemical and bio-based polymers.

31 **Introduction and updated taxonomical overview on marine OHCB**

32 Some 15 years ago we reported on an ecophysiologically unusual group of marine
33 microorganisms, termed 'obligate hydrocarbonoclastic (oil-degrading) bacteria (OHCB)
34 [1••]. One of their most distinct metabolic peculiarities is their ability for utilizing almost
35 exclusively hydrocarbons as a sole source of energy and carbon (a metabolic trait, called
36 the "OHCB paradigm" [1••]). Here, we need to note that genetic loci of this physiological trait

37 are relatively easily transmissible between bacteria, which is known for decades e.g. for
38 alkane hydroxylases/monooxygenases and P450 (CYP153) cytochromes
39 (<https://pubmed.ncbi.nlm.nih.gov/11390693/>) or for a half-century, aromatic
40 monooxygenases (TOL papers <https://pubmed.ncbi.nlm.nih.gov/4418209/>). Importantly,
41 the phylogenies of these enzymes are uncoupled with the taxonomic placement of
42 producing organisms (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4021335/>) therefore
43 the term OHCB cannot be strictly attributed to a particular taxon.

44
45 The ecological importance of OHCB in the biological removal of petroleum hydrocarbons
46 from various polluted marine environments around the world has currently been evidenced
47 during last dramatic oil spill disasters, such as aforementioned blowout of the Deepwater
48 Horizon (DWH) oil rig (see references below). Since the discovery of the first obligate marine
49 hydrocarbonoclastic bacteria being as only members of the class *Gammaproteobacteria*,
50 their taxonomic diversity has not much been changed at the level of higher taxa. To date, of
51 the eleven recognized genera, homing the OHCB species, only the genus *Planomicrobium*
52 includes the species *P. alkanoclasticus* [2], order *Firmicutes* within the class *Bacilli*, has
53 been added to the list. The rest of the well-established and novel OHCBs are the members
54 of the class *Gammaproteobacteria* and are subdivided into four orders: *Cellvibrionales*
55 (*Porticoccus* [3•]); *Nevskiales* (*Algiphilus* [4•] and *Polycyclovorans* [5•]); *Oceanospirillales*
56 (*Alcanivorax* [6], *Neptunomonas* [7], *Oleibacter* [8], *Oleiphilus* [9], *Oleispira* [10] and
57 *Thalassolituus* [11]) and *Thiothrichales* (*Cycloclasticus* [12]).

58 Initially characterized as highly specialized hydrocarbonoclastic bacteria that degrade either
59 aliphatic (*Alcanivorax*, *Oleibacter*, *Oleiphilus*, *Oleispira*, *Thalassolituus*) or aromatic
60 (*Cycloclasticus*, *Neptunomonas*) hydrocarbons [1•,13-15], OHCB possess a higher
61 metabolic versatility than previously assumed. For example, in addition to the type species
62 of the genus, *A. borkumensis* SK2^T, the genus *Alcanivorax*, currently includes 14 further

species with validly published names (<https://lpsn.dsmz.de/genus/alcanivorax>), some of which exhibit genome sizes much larger than that in the strain SK2^T and, in turn, utilize a larger range of growth substrates. More specifically, some *Alcanivorax* strains were found to be able to degrade simple sugars such as arabinose and glucose [16] and simple aromatic compounds, such as benzene, chlorobenzene, toluene [17] expanding the metabolic capability of this group of organisms, historically known as narrow specialists in degradation of aliphatic, branched hydrocarbons (including isoprenoids) and cycloalkanes [1•,6]. In concordance, some mussel and sponge symbionts isolated from deep-sea gas and oil seeps [18] were classified as the members of the genus *Cycloclasticus*, the genus of well-established primarily degraders of (poly)aromatic hydrocarbons in marine environments. They were found to be similar to free-living *Cycloclasticus* that bloomed during the DWH oil spill [19-24 and references therein]. Both these groups of bacteria were capable of degradation of short-chain alkanes ethane, propane and butane [18,19], thus highlighting the expanded role of these keystone species in the degradation of hydrocarbons released into marine environment. Of special attention are the recently uncovered novel OHCBs (*Algiphilus aromaticivorans*, *Polycyclovorans algicola*, *Porticoccus hydrocarbonoclasticus*) that represent novel genera and species and that were isolated from eukaryotic phytoplankton [3•,4•,5•]. All these isolates can use wide range of hydrocarbons, from aliphatic (C₁₀ – C₁₆) and branched (phytane, pristane) to mono- (benzene, toluene, p-xylene, biphenyl) and polyaromatic hydrocarbons (naphthalene, anthracene, phenanthrene, pyrene, fluorene) as sole carbon sources for growth [14,15].

84

85 **Marine OHCBs and the phycosphere**

86 Marine OHCBs appear to be confined to the marine environment, where they tend to be
87 highly enriched in petroleum-contaminated areas. However, in addition to polluted sites,

100 As highlighted in recent reviews, there are several biotic as well as abiotic sources from
101 which hydrocarbons, albeit in small quantities, can enter the ocean and sustain the
102 emergence of OHCBs in both pristine and remote areas [13,26]. One of the ubiquitous
103 sources of biotic hydrocarbons is chlorophyll A, the central pigment of photosynthesis, which
104 is an aromatic porphyrin ring with side chain of diterpene hydrocarbon (phytol) and is
105 accounting for 0.3% to 5% of the dry weight of microalgal and cyanobacterial cell [26]. It
106 should be noted that, as it has been known for a long time, many microalgae and especially
107 cyanobacteria are capable of producing significant quantities of various types of
108 hydrocarbons, including long-chain alkanes [27,28•,29•,30,31•], with estimated global flux
109 of de-novo produced hydrocarbons exceeding fossil petroleum inputs into ocean by 100-
110 500-fold [32•,33]. Therefore, it does not seem accidental that representatives of
111 *Alcanivorax*, *Thalassospira*, *Oleibacter* as well as new genera and species of OHCBs
112 (*Polycyclovorans*, *Algiphilus* and *Porticoccus hydrocarbonoclasticus*) have been commonly
113 reported to be associated with many species of phytoplankton (diatoms, dinoflagellates,
114 coccolithophores) [13,26,34-38]. In this regard, the global distribution of OHCBs in the
115 oceans can be explained by their conquest of still poorly studied and underexploited
116 biotopes - the cell surface or phycosphere [26] of marine phytoplankton. The association of
117 some OHCB taxa with these primary producers raises important questions regarding their
118 ecology and their contribution to ocean wellness. Oil pollution and biogenic production of
119 hydrocarbons by phytoplankton are corresponding aspects of the long- and short-term
120 hydrocarbon cycles in the oceans, in which OHCB communities, both phycosphere-
121 associated and free-living in seawater environments, play a significant role.

122 **Marine OHCBs and the plastisphere**

123 In addition to being very important for the sustainability of ocean ecosystems on a planetary
124 scale in respect to the hydrocarbon degradation, OHCB appear to play an additional and yet

underestimated role as an important part of the marine '*plastisphere*', the newly recognized microbial assemblages colonizing and modifying plastics released in the ocean [39-44]. Petroleum-derived synthetic plastics, including low- and high-density polyethylene (LDPE and HDPE), polystyrene (PS), polypropylene (PP), polyvinyl chloride (PVC), polyurethane (PUR), and polyethylene terephthalate (PET), are currently major pollutants of marine environments across the globe. It has recently been estimated that between 4.8 and 12.7 million Metric Tons (MTs) of plastic have entered the ocean annually over the past decade, largely due to improper land-based waste management [45]. According to the Plastics-Europe, the global yield of plastics reached 348 million MTs in 2018 [46]. Without major interventions in waste generation and recycling, it was predicted that by 2025, the cumulative amount of mismanaged plastic wastes entering the ocean could reach almost 250 million MTs [45,47]. On the other hand, the production of biodegradable bioplastics (BBPs) has been gradually increasing, as they are considered as promising alternatives to common petrochemistry-based polymers. In 2018, over 2 million MTs of BBPs was produced worldwide with the major BBP types including poly(lactic acid) (PLA, ~25%), poly(butylene succinate) (PBS, ~11%), poly(butylene adipate terephthalate) (PBAT, ~12%), polyhydroxyalkanoates (PHA, ~6%), and starch blends (~44%) [46]. The number of scientific studies focused on BBPs alone has been rapidly increasing with almost 18,000 publications in PubMed as of March 2021. However, BBPs polymers are recalcitrant in the marine environment, as exemplified by Napper and Thompson [48] and represent an emerging group of pollutants and, at the same time, a new substratum for microbial colonization. Recent studies have showed that the current mass of ocean plastic is in the concentration range from between 0.1-1.0 particles m⁻³ in the water column, and from 10³-10⁴ particles m⁻³ in sediments [49] and only 1% plastics entering the marine environment are observed floating at sea [41,50]. This observation is the opposite of the fact that over 65.5% of the

150 world's plastic is represented by lightweight polymers with a lower density than seawater
151 [51].

152 The interest to marine *plastisphere* has recently stimulated numerous studies, which
153 generated a breadth of data [40,52•,53]. Similar to the biofouling that occurs with any inert
154 solid material entering the sea, the plastisphere develops rapidly on the surface of plastic
155 debris when nutrients, carbon sources and energy are available. These determinants, as
156 well as salinity, temperature and solar irradiation, are more likely to influence the
157 development of a particular *plastisphere* [41], which may significantly differ in microbial
158 composition compared to free-living or particle-associated microbial populations in the
159 surrounding water [54. In addition, the type, size and status of the plastic itself (e.g. virgin or
160 weathered) also affects the formation of the *plastisphere*, composition of which is to some
161 extent characteristic for particular type of plastic [55••,56,57].

162 Given that common plastics are derived from fossil fuels, and especially, when oligomers
163 and plastic additives (plasticizers) are available as a source of carbon and energy, one
164 would expect that *plastisphere* should be enriched with OHCBs. The high hydrophobicity of
165 petro-sourced polyolefins (PE, PP and PS) as well as PET, PVC, and PUR creates a strong
166 interface when these materials are immersed in seawater, hindering attachment of the
167 majority of planktonic hydrophilic microbiota. However, these surfaces can be quickly
168 covered by marine microorganisms, possessing hydrophobic outer membrane. Worth to
169 mention, the regulation of the hydrophobicity of outer membranes from neat and hydrophilic
170 to indented and hydrophobic has been documented in a number of studies on marine
171 OHCBs [6,58-61]. Following these expectations, we have attempted to validate the role of
172 OHCBs in light of their abilities to both primary colonization (pioneering) and decomposition,
173 at least partially, different types of marine plastic debris. A more targeted objective was to

174 elucidate the specific role of OHCBs in the enzymatic (hydrolytic or redox) modification of
175 the original polymer structure.

176 **Marine OHCBs play important role in primo-colonization of plastic.**

177 Although there are many studies on the microbial composition of plastic biofilms, few studies
178 have been devoted to detailed analysis of the successive phases of plastisphere's
179 development [52,53]. Like biofouling [62,63], this process can be divided into three
180 sequential events: (i) "primo-colonization" phase, which implies the colonization of pioneer
181 bacteria on the plastic surface, which form the first layer of the original biofilm; (ii) a "growth
182 phase" that promotes irreversible attachment through formation of extracellular matrix such
183 as pili, adhesion proteins and exopolymeric substrate matrix (EPS) produced by both
184 primary and secondary species; (iii) the "maturation phase" occurs through diverse,
185 competitive or synergistic interactions between cells with further recruitment or loss of
186 species [56,64]. The role of OHCBs as key colonizers was first demonstrated by
187 investigating the different phases of the colonization of polyolefin-based plastics, namely
188 virgin low-density polyethylene (LDPE), PE with prooxidant (OXO), and artificially aged OXO
189 (AA-OXO) [56]. Succession of plastisphere formed on plastic pieces was monitored for 45
190 days, with all three developmental phases observed. Notably, OHCBs belonging to genera
191 *Alcanivorax*, *Oleiphilus* and *Thalassolituus* were especially abundant in the plastisphere
192 during the primo-colonization phase and tended to decrease thereafter [56]. It is very likely,
193 that LDPE attracts marine OHCBs independently of its status, since primo-colonization by
194 these organisms was also observed on the weathered LDPE [65]. Similar to the research's
195 data of Dussud et al. [56], the enrichment of a prominent OHCB, *Oleiphilus messinensis*, on
196 both non-weathered and weathered LDPE only occurred at early stages of colonization (i.e.,
197 after 2 days of incubation in coastal marine water [Mallorca, Spain]; where they represented
198 3.7% and 5.8% of relative abundance, respectively, vs. 0.6% on glass controls) [65]. To

199 some extent, a similar observation was obtained during colonization experiments conducted
200 with virgin high-density polyethylene (HDPE) microbeads (1-4 μm ; 0.96 g cm^{-3}) [66].
201 Incubation of seawater with HDPE for 108 hours stimulated the activity and enrichment of
202 six OTUs, two of which were closely related to OHCBs of the genera *Alcanivorax* and
203 *Oleispira*. The authors hypothesized that, due to an increase in oxygen consumption during
204 incubation with HDPE as compared to control, these specific organisms were uniquely
205 adapted to use organic carbon from virgin HDPE microbeads, likely plastic additives and /or
206 plasticizers, as carbon and energy sources, pointing at their potential for the bioremediation
207 of this type of plastic [66]. The plastisphere succession studied during colonization of three
208 types of polyethylene terephthalate (virgin and weathered PET power [particles <300 μm]
209 and amorphous PET films [250 μm thickness]) for 42 days led to slightly different results
210 [56]. In more details, members of *Piscirickettsiaceae*, the family accomodating the genus
211 *Cycloclasticus*, were found in significant numbers (relative abundance 4.35%) during all
212 phases of plastisphere development on all type of plastic studied. Two different amplified
213 sequence variants (ASV), ASV8 and ASV15 affiliated to *Alcanivoraceae* reached their
214 maximum abundance of 15% and 2.35%, respectively, but at later stages of the plastisphere
215 formation. Notably, while ASV8 was visualized only on virgin PET powder and amorphous
216 PET film, ASV15 was found to grow on all type of plastic studied. The joint participation of
217 *Alcanivorax* and *Cycloclasticus* species in formation of a mature biofilm on the surface of
218 PET films was confirmed in a separated study of Denaro et al. [67]. Interestingly, ATR-FTIR
219 and SEM analyses revealed that most significant alterations (formation of small cracks and
220 cavities) of the surface of PET films was mainly caused by the bacterial consortia enriched
221 on either tetradecane or diesel [67].

222 **Plastic-degrading enzymes from HCB.** As indicated above, marine OHCBs are likely to
223 survive in pristine environments through the use of alkanes secreted either by marine

224 cyanobacteria [30,31•,32••] or other eukaryotic primary producers [69]. Additionally, they
225 appear to have a strong hydrolytic capacity towards aliphatic polyesters, both naturally
226 occurring (e.g. polyhydroxyalkanoates [PHA]) and industrially produced (e.g. polybutylene
227 succinate [PBS], polycaprolactone [PCL], polyethylene succinate [PES] and polylactic acid
228 [PLA]) [70-72•]. Nowadays these aliphatic polyesters, referred above as BBPs, still account
229 for a small fraction of the global polymer market, but fortunately, consumer demand for these
230 “green” alternatives to traditional non-biodegradable petro-based materials is steadily
231 growing [73,74]. Although the molecular mechanisms underlying the ability of OHCB strains
232 to degrade such polyesters are currently unknown, genomes of OHCBs encode a large array
233 of enzyme candidates that potentially can be active on BBPs. Indeed, numerous enzymes
234 from *Alcanivorax*, *Oleispira*, and *Cycloclasticus* spp., either cloned from their genomes or
235 recovered from metagenomic DNA fragments in crude oil enrichments and attributed to
236 these organisms, were active on polyesters, including PLA, PBS, PHB and PET oligomers
237 [71,72•,75••-79•]. In particular, Tchigvintsev et al. [76•] and Hajighasemi et al. [77] reported
238 three ester hydrolases from the crude oil-seawater enrichments-derived metagenome, two
239 of which were 100% identical to proteins encoded by *A. borkumensis* SK2^T genome, one of
240 which, ABO2449, was hydrolyzing poly-(DL-lactic acid) polyesters. Notably, this activity of
241 ABO2449 was found in both emulsified and solid PLA, with the capacity to degrade solid
242 material being of high industrial interest. This polyester hydrolase had a highest activity at
243 30–37 °C and retained 32% of the maximum activity at 4 °C, suggesting that its cold-
244 adapted nature [80] ([Coscolin et al., 2018](#)). Search for novel and highly active polyesterases
245 from marine metagenomes has led to the discovery of the esterase GEN0105, which
246 hydrolyzes PCL, PLA as well as bis(benzoyloxyethyl)-terephthalate [78•]. Phylogenetic
247 analysis showed that this Family IV polyesterase, was closely related to the alpha/beta
248 hydrolases CUR46463 and WP_055099617 from *Alcanivorax xenomutans*. The recent
249 study on *Alcanivorax* sp. 24 isolated from marine plastic debris, which is able to degrade

250 BBP polyesters (PBS, PCL, PES and PHA), has also demonstrated its capacity of bis(2-
251 hydroxyethyl)-terephthalate (BHET) degradation. Further analysis of abundantly secreted
252 ALC24_4107 depolymerase revealed its promiscuous hydrolytic activity towards aliphatic
253 polyesters, which makes it possible to biodegrade both natural and synthetic polyester
254 plastics of anthropogenic origin [72•].

255 As pointed out elsewhere, many polyesters are of an unnatural origin, exhibit a low
256 bioavailability and are exposed to the microorganisms in their natural environment for a
257 limited period of time, which may be a reason why the evolution of enzymatic pathways for
258 plastic degradation *in situ* is a relatively slow process [41•,81]. Accordingly, the ubiquity of
259 polyester-utilising enzymes, e.g. homologs of PETase from *Idionella* [82••] in marine
260 bacteria and their communities, is correspondingly low, as reflected by the sequencing data
261 analysis [83•,84]. However, it is also known that some families of esterases exhibit an
262 enormous substrate promiscuity and some can utilize, e.g. 74 from 96 natural and synthetic
263 ester substrates tested [85••]. This also suggests that predictions of enzymatic activities
264 from genomic data may overlook enzymes that are readily available in nature and capable
265 of polyester degradation and points at importance of functional enzyme discovery and
266 experimental activity validation, e.g. proteins with domains of unknown function (DUF) or
267 “unknowns”.

268 **Conclusion and outlook**

269 The situation with the biodegradation of both natural and synthetic polyesters looks more
270 optimistic compared to the bioconversion of non-hydrolysable polyolefins of petrochemical
271 origin, such as polyethylene, polypropylene and polystyrene, which is hampered by the high
272 redox potential required to cleave carbon-carbon bonds. However, the ability of OHCB to
273 form thick biofilms specifically on LDPE, and break down this petroleum-based plastic has
274 recently been demonstrated [57]. As mentioned above, bacteria of the family
275 *Alcanivoracaceae* are also able to colonize the HDPE microbeads and, probably, use
276 organic carbon from this type of plastic [66] (Figure 2).

277



280

281

282

283

284

285

286

287

288

289

290

292 Only four types of enzymes (manganese and soybean peroxidases, laccase and alkane
293 hydroxylases [monooxygenases]) have exhibited the PE degradation activities [90]. Since
294 all OHCBs possess a large repertoire of enzymes of the latter type (AlkB, AlmA and P450
295 monooxygenases), their participation in degradation of low molecular-weight PE in marine
296 environment could be foreseen. It is worth to mention that a recent quantum mechanics
297 study predicting the catalytic mechanism of P450 monooxygenase suggested that
298 oxygenase-induced free radical transitions may cause the cleavage of the carbon-carbon
299 backbone of polyolefins (PE and PS) [87]. While these studies suggest a fundamental
300 possibility that the biodegradation processes of PE or PS, which is at least partially
301 performed by OHCB, are feasible in principle (Figure 2), more effort is needed to
302 characterize biochemical functions of corresponding enzyme candidates to help predicting
303 the plastics biodegradation pathways and enable the engineering of microbial catalysts for
304 plastics biodegradation or recycling [92•].

305 With the prevalence of plastic marine debris and the continued growth in plastic production,
306 the impact of plastic on marine ecosystems is likely to intensify. Currently, the possibilities
307 of microbial communities developing on the surface of plastic (marine plastisphere) to
308 change the structure and depolymerize both natural and synthetic polyesters are being
309 actively studied. We expect that OHCBs as part of the plastisphere – by accelerating the
310 characterization of their genetic circuits and repertoires – to make a significant contribution
311 to the biodegradation of plastics in the marine ecosystems.

312 **Conflict of interest statement**

313 The authors declare no conflict of interests.

314 **Acknowledgement**

315 This work was supported by research funds from the Italian Ministry of University and
316 Research (MUR), PRIN 2017, code: 201728ZA49_002 and by the European Commission's
317 Horizon Work Program H2020-FNR-2020 under FUTURENZYME Project Contract
318 (101000327).

319

320 **CRedit author statement**

321 **Michail M. Yakimov:** Conceptualization, Methodology, Writing- Original draft preparation.

322 **Rafael Bargiela:** Software, Data curation, Visualization. **Peter N. Golyshin:**

323 Conceptualization, Writing- Reviewing and Editing.

324

325

326 **References and recommended reading**

327 Paper of particular interest, published within the period of review, have been highlighted as:

328

329 • of special interest

330 •• of outstanding interest

331

332 1. Yakimov MM, Timmis KN, Golyshin PN: **Obligate oil-degrading marine bacteria.** *Curr Opin*
333 *Biotechnol* 2007, **18**:257-266.

334 ••One of the first review describing the physiologically novel group of hydrocarbon-degrading
335 marine bacteria

336

337 2. Engelhardt MA, Daly K, Swannell RPJ, Head IM: **Isolation and characterization of a novel**
338 **hydrocarbon-degrading, Gram-positive bacterium, isolated from intertidal beach sediment,**
339 **and description of *Planococcus alkanoclasticus* sp. nov.** *J Appl Microbiol* 2001, **90**:237-247.

340 3. Gutierrez T, Nichols PD, Whitman WB, Aitken MD: ***Porticoccus hydrocarbonoclasticus* sp.**
341 **nov., an aromatic hydrocarbon-degrading bacterium identified in laboratory cultures of**
342 **marine phytoplankton.** *Appl Environ Microbiol* 2012, **78**:628-637

343 • An important report introducing a novel member of OHCB group.

344 4. Gutierrez T, Green DH, Whitman WB, Nichols PD, Semple KT, Aitken MD: ***Algiphilus***
345 ***aromaticivorans* gen. nov., sp. nov., an aromatic hydrocarbon-degrading bacterium isolated**
346 **from a culture of the marine *dinoflagellate* *Lingulodinium polyedrum*, and proposal of**
347 ***Algiphilaceae* fam. nov.** *Int J Syst Evol Microbiol* 2012, **62**:2743-2749.

348 •An important report on a novel member of OHCB group.

349 5. Gutierrez T, Green DH, Nichols PD, Whitman WB, Semple KT, Aitken MD: ***Polycyclovorans***
350 ***algicola* gen. nov., sp. nov., an aromatic-hydrocarbon-degrading marine bacterium found**
351 **associated with laboratory cultures of marine phytoplankton.** *Appl Environ Microbiol* 2013,
352 **79**:205-214.

353 • An important report describing a novel member of OHCB group.

354 6. Yakimov MM, Golyshin PN, Lang S, Moore ER, Abraham WR, Lünsdorf H, Timmis KN:
355 ***Alcanivorax borkumensis* gen. nov., sp. nov., a new, hydrocarbon-degrading and**
356 **surfactant-producing marine bacterium.** *Int J Syst Bacteriol* 1998, **48**:339-348.

357 7. Hedlund BP, Geiselbrecht AD, Bair TJ, Staley JT: **Polycyclic aromatic hydrocarbon**
358 **degradation by a new marine bacterium, *Neptunomonas naphthovorans* gen. nov., sp. nov.**
359 *Appl Environ Microbiol* 1999, **65**:251-259

360 8. Teramoto M, Ohuchi M, Hatmanti A, Darmayati Y, Widyastuti Y, Harayama S, Fukunaga Y.
361 ***Oleibacter marinus* gen. nov., sp. nov., a bacterium that degrades petroleum aliphatic**
362 **hydro- carbons in a tropical marine environment.** *Int J Syst Evol Microbiol* 2011, **61**:375-380

363 9. Golyshin PN, Chernikova TN, Abraham WR, Lunsdorf H, Timmis KN, Yakimov MM:
364 ***Oleiphilaceae* fam. nov., to include *Oleiphilus messinensis* gen. nov., sp. nov., a novel**
365 **marine bacterium that obligately utilizes hydrocarbons.** *Int J Syst Evol Microbiol* 2002, **52**:901-
366 911

367 10. Yakimov MM, Giuliano L, Gentile G, Crisafi E, Chernikova TN, Abraham WR, Lünsdorf H,
368 Timmis KN, Golyshin PN: ***Oleispira antarctica* gen. nov., sp. nov., a novel**
369 **hydrocarbonoclastic marine bacterium isolated from Antarctic coastal sea water.** *Int J Syst*
370 *Evol Microbiol* 2003, **53**:779-785.

371 11. Yakimov MM, Giuliano L, Denaro R, Crisafi E, Chernikova TN, Abraham WR, Luensdorf H,
372 Timmis KN, Golyshin PN: ***Thalassolituus oleivorans* gen. nov., sp. nov., a novel marine**
373 **bacterium that obligately utilizes hydrocarbons.** *Int J Syst Evol Microbiol* 2004, **54**:141-148

374 12. Dyksterhouse SE, Gray JP, Herwig RP, Lara JC, Staley JT: ***Cycloclasticus pugetii* gen.**
375 **nov., sp. nov., an aromatic hydrocarbon- degrading bacterium from marine sediments.** *Int J*
376 *Syst Bacteriol* 1995, **45**:116-123

377 13. McGenity TJ, Folwell BD, McKew BA, Sanni GO: **Marine crude-oil biodegradation: a central**
378 **role for interspecies interactions.** *Aquat Biosyst* 2012, **8**:10.

379 14. Gutierrez T: **Marine, aerobic hydrocarbon-degrading gammaproteobacteria: overview.** In
380 *Taxonomy, Genomics and Ecophysiology of Hydrocarbon-Degrading Microbes*. Edited by
381 McGenity TJ. Springer, Cham, 2017:1-10.

382 15. Gutierrez T: **Aerobic hydrocarbon-degrading Gammaproteobacteria: *Porticoccus*.** In
383 *Taxonomy, Genomics and Ecophysiology of Hydrocarbon-Degrading Microbes*. Edited by
384 McGenity TJ. Springer, Cham, 2019:181-189.

385 16. Yakimov MM, Golyshin PN, Crisafi F, Denaro R, Giuliano L: **Marine, aerobic hydrocarbon-**
386 **degrading Gammaproteobacteria: the family *Alcanivoracaceae*.** In *Taxonomy, Genomics and*
387 *Ecophysiology of Hydrocarbon-Degrading Microbes*. Edited by McGenity TJ. Springer, Cham,
388 2019:167-179.

389 17. Hassan HA, Rizk NMH, Hefnawy MA, Awad AM: **Isolation and characterization of halophilic**
390 **aromatic and chloroaromatic degrader from Wadi El-Natrun Soda lakes.** *Life Sci J* 2012, **9**:
391 1565-1570.

392 18. Rubin-Blum M, Antony CP, Borowski C, Sayavedra L, Pape T, Sahling H, Bohrmann G, Kleiner
393 M, Redmond MC, Valentine DL, Dubilier N: **Short-chain alkanes fuel mussel and sponge**
394 ***Cycloclasticus* symbionts from deep-sea gas and oil seeps.** *Nat Microbiol* 2017, **2**:17093.

395 19. Valentine DL, Kessler JD, Redmond MC, Mendes SD, Heintz MB, Farwell C, Hu L, Kinnaman
396 FS, Yvon-Lewis S, Du M, Chan EW, Tigreros FG, Villanueva CJ: **Propane respiration jump**
397 **starts microbial response to a deep oil spill.** *Science* 2010, **330**:208-211.

398 20. Gutierrez T, Singleton DR, Berry D, Yang T, Aitken MD, Teske A: **Hydrocarbon-degrading**
399 **bacteria enriched by the *Deepwater Horizon* oil spill identified by cultivation and DNA-SIP.**
400 *ISME J* 2013, **7**:2091-2104.

401 21. Kleindienst S, Grim S, Sogin M, Bracco A, Crespo-Medina M, Joye SB: **Diverse, rare**
402 **microbial taxa responded to the *Deepwater Horizon* deep-sea hydrocarbon plume.** *ISME J*
403 2016, **10**:400-415.

404 22. Dombrowski N, Donaho J, Gutierrez T, Seitz KW, Teske AP, Baker BJ: **Reconstructing**
405 **metabolic pathways of hydrocarbon-degrading bacteria from the *Deepwater Horizon* oil**
406 **spill.** *Nat Microbiol* 2016, **1**:16057.

415 23. Yang T, Speare K, McKay L, MacGregor BJ, Joye SB, Teske A: **Distinct bacterial**
416 **communities in surficial seafloor sediments following the 2010 *Deepwater Horizon* blowout.**
417 *Front Microbiol* 2016, **7**:1384.

418

419 24. Gutierrez T, Morris G, Ellis D, Bowler B, Jones M, Salek K, Mulloy B, Teske A: **Hydrocarbon-**
420 **degradation and MOS-formation capabilities of the dominant bacteria enriched in sea**
421 **surface oil slicks during the *Deepwater Horizon* oil spill.** *Mar Pollut Bull* 2018, **135**:205-215.

422

423 25. Padgham M, Rudis B, Lovelace R, Salmon M: **osmdata.** *J Open Source Softw* 2017, **2**:315.

424

425 26. Gutierrez T: **Occurrence and roles of the obligate hydrocarbonoclastic bacteria in the**
426 **ocean when there is no obvious hydrocarbon contamination.** In *Taxonomy, Genomics and*
427 *Ecophysiology of Hydrocarbon-Degrading Microbes*. Edited by McGenity TJ. Springer, Cham;
428 2019.

429

430 27. Han J, McCarthy ED, Hoeven WV, Calvin M, Bradley WH: **Organic geochemical studies II. A**
431 **preliminary report on the distribution of aliphatic hydrocarbons in algae, in bacteria, and in**
432 **recent lake sediment.** *Proc Natl Acad Sci U S A* 1968, **59**:29-33.

433

434 28. Han J, Calvin M: **Hydrocarbon distribution of algae and bacteria, and microbiological**
435 **activity in sediments.** *Proc Natl Acad Sci U S A* 1969, **64**:436-43.

436 ••One of the first reports on very significant yields of hydrocarbons (primarily, medium-chain
437 alkanes) production in cyanobacteria ("blue-green algae")

438

439 29. Schirmer A, Rude MA, Li X, Popova E, del Cardayre SB: **Microbial biosynthesis of alkanes.**
440 *Science* 2010, **329**:559-562.

441 •An important report on pathways for cyanobacterial biosynthesis of alkanes.

442

443 30. Coates RC, Podell S, Korobeynikov A, Lapidus A, Pevzner P: **Characterization of**
444 **cyanobacterial hydrocarbon composition and distribution of biosynthetic pathways.** *PLoS*
ONE 2014, **9**:85140.

445

446 31. Lea-Smith DJ, Biller SJ, Davey MP, Cotton CAR, Sepulveda BMP, Turchyn AV, Scanlan DJ,
447 Smith AG, Chisholm SW, Howe CJ: **Contribution of cyanobacterial alkane production to the**
ocean hydrocarbon cycle. *Proc Natl Acad Sci USA* 2015, **112**:13591-13596.

448 •A series of laboratory microcosm experiments with alkane producing cyanobacteria and *A.*
449 *borkumensis* SK2 suggest the OHCB can sustain its growth on pentadecane from photosynthetic
450 alkane producer. It was predicted that the short-term microbial hydrocarbon loop may globally be
451 of a great scale.

452

453 32. Love CR, Arrington EC, Gosselin KM, Reddy CM, Van Mooy BAS, Nelson RK, Valentine DL:
454 **Microbial production and consumption of hydrocarbons in the global ocean.** *Nat Microbio*
2021, **6**:489-498.

455 ••In situ hydrocarbon production measurements and scale-up modelling suggested the scale of the
456 "short-term" (or "cryptic") hydrocarbon cycle outweighs, at least hundredfold, the petroleum input
457 into the global ocean.

458

459 33. Harada N, Hirose Y, Chihong S, Kurita H, Sato M, Onodera J, Murata K, Itoh F: **A novel**
460 **characteristic of a phytoplankton as a potential source of straight-chain alkanes.** *Sci Rep*
2021, **11**:14190.

34. Green DH, Llewellyn LE, Negri AP, Blackburn SI, Bolch CJS: **Phylogenetic and functional diversity of the cultivable bacterial community associated with the paralytic shellfish poisoning dinoflagellate *Gymnodinium catenatum***. *FEMS Microbiol Ecol* 2004, **47**:345-357.

35. Green DH, Bowman JP, Smith EA, Gutierrez T, Bolch CJS: ***Marinobacter algicola* sp. nov., isolated from laboratory cultures of paralytic shellfish toxin-producing dinoflagellates**. *Int J Syst Evol Microbiol* 2006, **56**:523-527.

36. Green DH, Echavarri-Bravo V, Brennan D, Hart MC: **Bacterial diversity associated with the coccolithophorid algae *Emiliania huxleyi* and *Coccolithus pelagicus* f. *braarudii***. *Biomed Res Int* 2015:194540.

37. Chernikova TN, Bargiela R, Toshchakov SV, Shivaraman V, Lunev EA, Yakimov MM, Thomas DN, Golyshin PN: **Hydrocarbon-degrading bacteria *Alcanivorax* and *Marinobacter* associated with microalgae *Pavlova lutheri* and *Nannochloropsis oculata***. *Front Microbiol* 2020; **11**:572931.

38. Thompson HF, Summers S, Yuecel R, Gutierrez T: **Hydrocarbon-degrading bacteria found tightly associated with the 50-70 µm cell-size population of eukaryotic phytoplankton in surface waters of a northeast Atlantic Region**. *Microorganisms* 2020, **8**:1955.

39. Amaral-Zettler LA, Zettler ER, Slikas B, Boyd GD, Melvin DW, Morrall CE, Proskurowski G, Mincer TJ: **The biogeography of the Plastisphere: implications for policy**. *Front Ecol Environ* 2015, **13**: 541-546

40. Amaral-Zettler LA, Zettler ER, Mincer TJ: **Ecology of the Plastisphere**. *Nat Rev Microbiol* 2020, **18**:139-151.

41. Oberbeckmann S, Labrenz M: **Marine microbial assemblages on microplastics: diversity, adaptation, and role in degradation**. *Annu Rev Mar Sci* 2020, **12**:1-24.

•A comprehensive critical analysis of a multitude of studies on marine “Plastisphere”

42. Ogonowski M, Motiei A, Ininbergs K, Hell E, Gerdes Z, Udekwu KI, Bacsik Z, Gorokhova E: **Evidence for selective bacterial community structuring on microplastics**. *Environ Microbiol* 2018, **20**:2796-2808.

43. Reisser J, Shaw J, Hallegraeff G, Proietti M, Barnes DKA, Thums M, Wilcox C, Hardesty BD, Pattiaratchi C: **Millimeter-sized marine plastics: a new pelagic habitat for microorganisms and invertebrates**. *PLoS One* 2014, **9**:e100289.

44. Rummel CD, Jahnke A, Gorokhova E, Kühnel D, Schmitt-Jansen M: **The Impacts of biofilm formation on the fate and potential effects of microplastic in the aquatic environment**. *Environ Sci Technol Lett* 2017, **4**:258-267.

45. Jambeck J, Geyer R, Wilcox C, Siegler T, Perryman M, Andrady A, Narayan R, Law K: **Marine pollution. Plastic waste inputs from land into the ocean**. *Science* 2015, **347**:768-771.

46. Plastics Europe: **Plastics - The Facts 2018: An Analysis of European Plastics Production, Demand and Waste Data (EB/OL)**. 2018, URL: <http://www.plasticseurope.org/>

47. Galgani L, Loiselle SA: **Plastic pollution impacts on marine carbon biogeochemistry** *Environ Pollut* 2021; **268**:115598 doi: 10.1016/j.envpol.2020.115598

504 48. Napper IE, Thompson RC: **Environmental deterioration of biodegradable, oxo-**
505 **biodegradable, compostable, and conventional plastic carrier bags in the sea, soil, and**
506 **open-air over a 3-year period.** *Environ Sci Technol* 2019, **53**:4775- 4783.

507 49. Erni-Cassola G, Zadjelovic V, Gibson MI, Christie-Oleza JA: **Distribution of plastic polymer**
508 **types in the marine environment; a meta-analysis.** *J Hazard Mater* 2019; **369**:691-698.

509 50. Van Sebillie E, Wilcox C, Lebreton L, Maximenko N, Hardesty BD, Van Franeker JA, Eriksen M,
510 Siegel D, Galgani F, Law KLA: **Global inventory of small floating plastic debris.** *Environ Res*
511 *Lett* 2015; **10**(12):1-11.

512 51. Geyer R, Jambeck JR, Law KL: **Production, use, and fate of all plastics ever made.** *Sci Adv*
513 2017; **3**:e1700782.

514 52. Wright RJ, Erni-Cassola G, Zadjelovic V, Latva M, Christie-Oleza J: **Marine plastic debris - a**
515 **new surface for microbial colonization.** *Environ Sci Technol* 2020, **54**:11657-72.
516 •A highly recommended critical overview of various aspects of colonization and degradation of
517 plastics in marine environments

518 53. Wright RJ, Langille MGI, Walker TR: **Food or just a free ride? A meta-analysis reveals the**
519 **global diversity of the Plastisphere.** *ISME J* 2021, **15**:789-806.

520 54. Yang Y, Liu W, Zhang Z, Grossart H-P, Gadd GM: **Microplastics provide new microbial**
521 **niches in aquatic environments.** *Appl Microbiol Biotechnol* 2020, **104**:6501-651.

522 55. Zettler ER, Mincer TJ, Amaral-Zettler LA: **Life in the “plastisphere”: microbial communities**
523 **on plastic marine debris.** *Environ Sci Technol* 2013, **47**:7137-7146.
524 ••An important study, which conceptualized the plastic surface as a distinct environment for
525 microbial colonization

526 56. Dussud C, Hudec C, George M, Fabre P, Higgs P, Bruzard S, Delort A-M, Eyheraguibel B,
527 Meistertzheim A-L, Jacquin J, Cheng J, Callac N, Odobel C, Rabouille S, Ghiglione J-F:
528 **Colonization of non-biodegradable and biodegradable plastics by marine microorganisms.**
529 *Front Microbiol* 2018; **9**:1571.

530 57. Delacuvellerie A, Cyriaque V, Gobert S, Benali S, Wattiez R. **The plastisphere in marine**
531 **ecosystem hosts potential specific microbial degraders including *Alcanivorax borkumensis***
532 **as a key player for the low-density polyethylene degradation.** *J Hazard Mat* 2019; **280**:1-
533 120899

534 58. Abraham WR, Meyer H, Yakimov M: **Novel glycine containing glucolipids from the alkane**
535 **using bacterium *Alcanivorax borkumensis*.** *Biochim Biophys* 1998; **1393**:57-62. doi:
536 10.1016/S0005-2760(98)00058-7

537 59. Manilla-Pérez E, Lange AB, Hetzler S, Steinbüchel A: **Occurrence, production, and export**
538 **of lipophilic compounds by hydrocarbonoclastic marine bacteria and their potential use to**
539 **produce bulk chemicals from hydrocarbons.** *Appl Microbiol Biotechnol* 2010, **86**:1693-1706.

540 60. Naether DJ, Slawtschew S, Stasik S, Engel M, Olzog M, Wick LY: **Adaptation of the**
541 **hydrocarbonoclastic bacterium *Alcanivorax borkumensis* SK2 to alkanes and toxic organic**
542 **compounds: a physiological and transcriptomic approach.** *Appl Environ Microbiol* 2013,
543 **79**:4282-4293.

553 61. Barbato M, Scoma A, Mapelli F, De Smet R, Banat IM, Daffonchio D, Boon N, Borin S:
554 **Hydrocarbonoclastic *Alcanivorax* isolates exhibit different physiological and expression**
555 **responses to n-dodecane.** *Front Microbiol* 2016, **7**:2056.
556

557 62. Dang H, Lovell CR: **Bacterial primary colonization and early succession on surfaces in**
558 **marine waters as determined by amplified rRNA gene restriction analysis and sequence**
559 **analysis of 16S rRNA genes.** *Appl Environ Microbiol* 2020, **66**:467-475.
560

561 63. Salta M, Wharton JA, Blache Y, Stokes KR, Briand JF: **Marine biofilms on artificial surfaces:**
562 **structure and dynamics.** *Environ Microbiol* 2013, **15**:2879-2893.
563

564 64. Lorite GS, Rodrigues CM, De Souza AA, Kranz C, Mizaikoff B, Cotta MA: **The role of**
565 **conditioning film formation and surface chemical changes on *Xylella fastidiosa* adhesion**
566 **and biofilm evolution.** *J Colloid Interface Sci* 2011, **359**:289-295.

567 65. Erni-Cassola G, Wright RJ, Gibson MI, Christie-Oleza JA: **Early colonization of weathered**
568 **polyethylene by distinct bacteria in marine coastal seawater.** *Microb Ecol* 2020, **79**:517-526.

569 66. Girard EB, Kaliwoda M, Schmahl WW, Wörheide G, Orsi WD: **Biodegradation of textile**
570 **waste by marine bacterial communities enhanced by light.** *Environ Microbiol Rep* 2021,
571 **12**:406-418.

572 67. Denaro R, Aulenta F, Crisafi F, Di Pippo F, Cruz Viggi C, Matturro B, Tomei P, Smedile F,
573 Martinelli A, Di Lisio V, Venezia C, Rossetti S: **Marine hydrocarbon-degrading bacteria**
574 **breakdown poly(ethylene terephthalate) (PET).** *Sci Total Environ* 2020, **749**:141608.
575

576 68. Sorigué D, Legeret B, Cuine S, Morales P, Mirabella B, Guedeney G, Li-Beisson Y, Jetter R,
577 Peltier G, Beisson F: **Microalgae synthesize hydrocarbons from long-chain fatty acids via a**
578 **light-dependent pathway.** *Plant Physiol* 2016, **171**:2393- 2405.

579 69. Sorigué D, Légeret B, Cuiné S, Blangy S, Moulin S, Billon E, Richaud P, Brugière S, Couté Y,
580 Nurizzo D, Müller P, Brettel K, Pignol D, Arnoux P, Li-Beisson Y, Peltier G, Beisson F: **An algal**
581 **photoenzyme converts fatty acids to hydrocarbons.** *Science* 2017, **357**:903-907.

582 70. Sekiguchi T, Saika A, Nomura K, Watanabe T, Watanabe T, Fujimoto Y, Enoki M, Sato T, Kato
583 C, Kanehiro H: **Biodegradation of aliphatic polyesters soaked in deep seawaters and**
584 **isolation of poly(ε-caprolactone)-degrading bacteria.** *Polym Degrad Stab* 2011, **96**:1397- 1403.
585

586 71. Zadjelovic V, Gibson MI, Dorador C, Christie-Oleza JA: **Genome of *Alcanivorax* sp. 24: a**
587 **hydrocarbon degrading bacterium isolated from marine plastic debris.** *Mar Genomics* 2019,
588 **49**:1-4.
589

590 72. Zadjelovic V, Chhun A, Quareshy M, Silvano E, Hernandez-Fernaund JR, Aguilo-Ferretjans MM,
591 Bosch R, Dorador C, Gibson MI, Christie-Oleza JA: **Beyond oil degradation: enzymatic**
592 **potential of *Alcanivorax* to degrade natural and synthetic polyesters.** *Environ Microbiol* 2020,
593 **22**:1356-1369.
594 •*Alcanivorax* sp. 24 isolated from marine plastic debris has a larger genome and a broader
595 substrate spectrum than the 'classical' OHCB strain SK2; it encodes an array of enzymes active on
596 a range of polyester substrates, including BBP
597

598 73. Tokiwa Y, Calabia BP, Ugwu CU, Aiba S: **Biodegradability of plastics.** *Int J Mol Sci* 2009,
599 **10**:3722-3742.
600

74. Aeschelmann F, Carus M: **Biobased building blocks and polymers in the world: capacities, production, and applications-status quo and trends towards 2020**. *Ind Biotechnol* 2015, **11**:154-159.

75. Kube M, Chernikova TN, Al-Ramahi Y, Beloqui A, Lopez-Cortez N, Guazzaroni ME, Heipieper HJ, Klages S, Kotsyurbenko OR, Langer I, Nechitaylo TY, Lunsdorf H, Fernandez M, Juarez S, Ciordia S, Singer A, Kagan O, Egorova O, Petit PA, Stogios P, Kim Y, Tchigvintsev A, Flick R, Denaro R, Genovese M, Albar JP, Reva ON, Martinez-Gomariz M, Tran H, Ferrer M, Savchenko A, Yakunin AF, Yakimov MM, Golyshina OV, Reinhardt R, Golyshin PN: **Genome sequence and functional genomic analysis of the oil-degrading bacterium *Oleispira antarctica***. *Nat Commun* 2013, **4**:2156.

•A comprehensive analysis of a cold-adapted OHCB important in deep cold and polar seas. Cold-adapted chaperonin interactome suggested proteins important for life in the cold. Dozens of enzymes (including carboxylesterases) have been cloned and characterized. Surprisingly, majority of proteins had mesophilic activity optima. The study has more than doubled the number of cold-adapted proteins in PDB.

76. Tchigvintsev A, Tran H, Popovic A, Kovacic F, Brown G, Flick R, Hajighasemi M, Egorova O, Somody JC, Tchigvintsev D, Khusnutdinova A, Chernikova TN, Golyshina OV, Yakimov MM, Savchenko A, Golyshin PN, Jaeger K-E, Yakunin AF: **The environment shapes microbial enzymes: five cold-active and salt-resistant carboxylesterases from marine metagenomes**. *Appl Microbiol Biotechnol* 2015, **99**:2165-2178.

•A number of esterases active on polyesters from crude oil enrichment metagenomes active on polyesters

77. Hajighasemi M, Nocek BP, Tchigvintsev A, Brown G, Flick R, Xu X, Cui H, Hai T, Joachimiak A, Golyshin PN, Savchenko A, Edwards EA, Yakunin AF: **Biochemical and structural insights into enzymatic depolymerization of polylactic acid and other polyesters by microbial carboxylesterases**. *Biomacromolecules* 2016, **17**:2027-2039.

78. Hajighasemi M, Tchigvintsev A, Nocek B, Flick R, Popovic A, Hai T, Khusnutdinova AN, Brown G, Xu X, Cui H, Anstett J, Chernikova TN, Bröls T, Le Paslier D, Yakimov MM, Joachimiak A, Golyshina OV, Savchenko A, Golyshin PN, Edwards EA, Yakunin AF: **Screening and characterization of novel polyesterases from environmental metagenomes with high hydrolytic activity against synthetic polyesters**. *Environ Sci Technol* 2018, **52**:12388-12401.

•Esterases including those from OHCB genomes or crude oil-enriched metagenomes were active on polyesters, PLA, PCL, PBSA and 3PET. The latter substrate was converted by four enzymes from OHCB *A. borkumensis* SK2.

79. Popovic A, Hai T, Tchigvintsev A, Hajighasemi M, Nocek B, Khusnutdinova AN, Brown G, Glinos J, Flick R, Skarina T, Chernikova TN, Yim V, Bröls T, Paslier DL, Yakimov MM, Joachimiak A, Ferrer M, Golyshina OV, Savchenko A, Golyshin PN, Yakunin AF: **Activity screening of environmental metagenomic libraries reveals novel carboxylesterase families**. *Sci Rep* 2017, **7**:44103.

•Screening of >1 M metagenomic clones led to a full characterization of >80 esterases active on polyesters, e.g. PLA and PCL. Study revealed active enzymes from new protein families, DUF3089 and “unknown”. Both were derived from hydrocarbon-degrading microbial communities: from naphthalene-spiked sample enriched in *Cycloclasticus* sp. (Messina harbour) and from crude oil-grown biofilm from the hull of the tanker M/T Haven sunken in 1991 off Genoa.

80. Coscolín C, Bargiela R, Martínez-Martínez M, Alonso S, Bollinger A, Thies S, Chernikova TN, Hai T, Golyshina OV, Jaeger K-E, Yakimov MM, Golyshin PN, Ferrer M: **Hydrocarbon-degrading microbes as sources of new biocatalysts**, In *Taxonomy, genomics and ecophysiology of hydrocarbon-degrading microbes*. Edited by McGenity TJ. Springer, Cham, 2018:353-373. https://doi.org/10.1007/978-3-030-14796-9_13

81. Krueger MC, Harms H, Schlosser D: **Prospects for microbiological solutions to environmental pollution with plastics.** *Appl Microbiol Biotechnol* 2015, **99**:8857-8874.

82. Yoshida S, Hiraga K, Takehana T, Taniguchi I, Yamaji H, Maeda Y, Toyohara K, Miyamoto K, Kimura Y, Oda K: **A bacterium that degrades and assimilates poly(ethylene terephthalate).** *Science* 2016, **351**:1196-99

••A landmark discovery of a PET-mineralising bacterium *Ideonella sakaiensis* with the two key enzymes for hydrolysis of PET and the reaction intermediate, mono(2-hydroxyethyl) terephthalic acid to produce easily biodegradable ethylene glycol and terephthalic acid.

83. Danso D, Schmeisser C, Chow J, Zimmermann W, Wei R, Leggewie C, Li X, Hazen T, Streit WR: **New insights into the function and global distribution of polyethylene terephthalate (PET)-degrading bacteria and enzymes in marine and terrestrial metagenomes.** *Appl Environ Microbiol* 2018, **84**: e02773-17.

•A global survey of marine and terrestrial metagenomes has identified >800 enzyme candidates predicted to hydrolyse PET, few enzymes were expressed and their activities confirmed. One of characterised PETases was derived from the OHCB, *Oleispira antarctica* RB-8.

84. Danso D, Chow J, Streit WR: **Plastics: environmental and biotechnological perspectives on microbial degradation.** *Appl Environ Microbiol* 2019, **85**:e01095-19.

85. Martínez-Martínez M, Coscolín C, Santiago G, Chow J, Stogios PJ, Bargiela R, Gertler C, Navarro- Fernández J, Bollinger A, Thies S, Méndez-García C, Popovic A, Brown G, Chernikova TN, García-Moyano A, Bjerga GEK, Pérez-García P, Hai T, Del Pozo MV, Stokke R, Steen IH, Cui H, Xu X, Nocek BP, Alcaide M, Distaso M, Mesa V, Peláez AI, Sánchez J, Buchholz PCF, Pleiss J, Fernández-Guerra A, Glöckner FO, Golyshina OV, Yakimov MM, Savchenko A, Jaeger KE, Yakunin AF, Streit WR, Golyshin PN, Guallar V, Ferrer M, The Inmare Consortium: **Determinants and prediction of esterase substrate promiscuity patterns.** *ACS Chem Biol* 2018, **13**:225-234.

•• Substrate profiling of approx. 150 carboxylesterases, including enzymes from OHCB, revealed a remarkable substrate promiscuity in enzymes of Family VI: some enzymes converted up to 74 of 96 tested ester substrates, including non-natural esters. Structural modelling suggested the “active site effective volume” as a main predictive marker for substrate promiscuity.

86. Ru J, Huo Y, Yang Y: Microbial degradation and valorization of plastic wastes. *Front Microbiol* 2020, **11**:442.

87. Xu J, Cui Z, Nie K, Cao H, Jiang M, Xu H, Tan T, Liu L: **A quantum mechanism study of the cc bond cleavage to predict the bio-catalytic polyethylene degradation.** *Front Microbiol* 2019, **10**:489.

88. Katoh K, Standley DM: **MAFFT multiple sequence alignment software version 7: improvements in performance and usability.** *Mol Biol Evol* 2013, **30**:772-780.

89. Capella-Gutiérrez S, Silla- Martínez JM, Gabaldón T: **trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses.** *Bioinformatics* 2009, **25**:1972-1973.

90. Team RDC: **R: A language and environment for statistical computing.** *R Foundation for Statistical Computing* 2010.

91. Paradis E, Schliep K: **ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R.** *Bioinformatics* 2019, **35**:526-528.

704 92. Wei R, Tiso T, Bertling J, O'Connor K, Blank LM, Bornscheuer UT: **Possibilities and**
705 **limitations of biotechnological plastic degradation and recycling.** *Nat Catal* 2020, **3**:867-871.
706 • A very thorough overview about perspectives and challenges of biotech approaches to for plastic
707 biodegradation and recycling.
708

709

710